

ПРИКЛАДНАЯ МАТЕМАТИКА

UDC 004.8
MSC 03E72

Union and meet of an infinite number of type-2 fuzzy sets*

O. V. Baskov

St. Petersburg State University, 7–9, Universitetskaya nab., St. Petersburg,
199034, Russian Federation

For citation: Baskov O. V. Union and meet of an infinite number of type-2 fuzzy sets. *Vestnik of Saint Petersburg University. Applied Mathematics. Computer Science. Control Processes*, 2021, vol. 17, iss. 2, pp. 108–119. <https://doi.org/10.21638/11701/spbu10.2021.201>

The article examines the infimum and supremum of an infinite number of fuzzy numbers. It is shown that familiar properties of these operations, which are valid for real numbers, may apply to fuzzy numbers only under certain conditions. A formula for computing the infimum and supremum of any set of fuzzy numbers is provided. Since the union and meet of type-2 fuzzy sets are defined via the infimum and supremum of fuzzy numbers, all the results obtained are applicable to these operations as well.

Keywords: type-2 fuzzy sets, union, meet, fuzzy numbers, infimum, supremum.

1. Introduction. Fuzzy sets introduced by L. A. Zadeh [1] have numerous applications in various fields of research due to their ability to deal with uncertain information. Usual crisp sets may be characterized by an indicator function that takes two values: 1 for elements belonging to a set, and 0 for all others. In fuzzy sets this function is called membership function, and it is permitted to take any value between 0 and 1. Higher values of the membership function correspond to higher degrees of certainty in whether an element should be part of a set. When the elements are the reals, fuzzy sets are called fuzzy numbers.

The next step in generalization of membership is allowing the values of the membership function themselves to be fuzzy [2]. So in type-2 fuzzy sets membership functions map elements to the fuzzy numbers. Some studies [3, 4] have found that increased fuzziness makes type-2 fuzzy sets better suited for certain tasks than type-1 fuzzy sets. Thus, it is undoubted that type-2 fuzzy sets will have many applications, so studying their properties is an important task.

M. Mizumoto and K. Tanaka [5] examined the set-theoretic operations on a finite number of type-2 fuzzy sets. In some areas, for instance, in an axiomatic approach to

* The work is supported by Russian Foundation for Basic Research (project N 20-07-00298-a).

© St. Petersburg State University, 2021

Pareto set reduction [6], the need arises to operate on infinite number of type-2 fuzzy sets. In this case the results from [5, 7] are not directly applicable. The aim of this paper is to determine the conditions under which the set-theoretic operations preserve their usual properties when applied to an infinite number of type-2 fuzzy sets. And since these operations are closely related to operations on the values of membership functions, i. e. on fuzzy numbers, we simultaneously study the properties of infimum and supremum of an infinite set of fuzzy numbers.

Operations on type-2 fuzzy sets are derived using Zadeh's extension principle. Such definitions are inconvenient for direct computations, so various algorithms and formulae exist to simplify performing set-theoretic operations on certain classes of type-2 fuzzy sets. One such formula was given by N. N. Karnik and J. M. Mendel [7] for computing join and meet of a finite number of type-2 fuzzy sets. It is applicable to a broad class of type-2 fuzzy sets, namely, the sets with strongly normal and convex values of membership functions. Our main result generalizes this formula to accept infinite, possibly uncountable number of type-2 fuzzy sets.

2. Preliminaries. Let X be the universal set of objects of any kind. A (*type-1*) *fuzzy set* A over X is a set of pairs $(x, \mu_A(x))$, where $x \in X$, $\mu_A(x) \in [0; 1]$. The number $\mu_A(x)$ is called a *degree of membership* of an element x in the fuzzy set A . The statement “ x certainly belongs to A ” corresponds to $\mu_A(x) = 1$, the assertion $x \notin A$ is written as $\mu_A(x) = 0$. Values of $\mu_A(x)$ between 0 and 1 represent uncertainty about whether x should be part of the set A or not: the higher the value $\mu_A(x)$, the more confident we are that $x \in A$. The function $\mu_A : X \mapsto [0; 1]$ is called the *membership function* of the fuzzy set A . Since a fuzzy set is fully determined by its membership function, we will often use them interchangeably.

Let A and B be two fuzzy sets over the universal set X . Their union $A \cup B$, intersection $A \cap B$, and complement \overline{A} are defined as follows: $\mu_{A \cup B}(x) = \max\{\mu_A(x); \mu_B(x)\}$, $\mu_{A \cap B}(x) = \min\{\mu_A(x); \mu_B(x)\}$, $\mu_{\overline{A}}(x) = 1 - \mu_A(x)$. It is said that $A \subseteq B$ if $\mu_A(x) \leq \mu_B(x)$ for every $x \in X$.

An α -cut of a fuzzy set μ_A is the crisp set $A_\alpha = \{x \in X : \mu_A(x) \geq \alpha\}$.

A *height* of a fuzzy set μ_A is $h_A = \sup_{x \in X} \mu_A(x)$. If $h_A = 1$, the fuzzy set μ_A is called *normal*. If furthermore $\mu_A(x) = 1$ for some $x \in X$, the fuzzy set μ_A is called *strongly normal*.

A fuzzy set μ_A over a convex set X is itself called *convex* if for any $\lambda \in (0; 1)$ and $x, y \in X$ it is true that $\mu_A(\lambda x + (1 - \lambda)y) \geq \min\{\mu_A(x); \mu_A(y)\}$.

A *fuzzy number* is a fuzzy set over the reals R . For fuzzy numbers the convexity condition may be rewritten in an easier form: a fuzzy number μ is convex if $\mu(y) \geq \min\{\mu(x); \mu(z)\}$ for any $x < y < z$.

A fuzzy number μ is *upper semicontinuous* if its α -cuts are closed for any $\alpha \in [0; 1]$. Observe that upper semicontinuous normal fuzzy numbers are necessarily strongly normal.

Operations on fuzzy numbers are derived using Zadeh's extension principle [2]: given some binary operation \circ on reals, one may define $\mu_{P \circ Q}(x) = \sup_{u \circ v = x} \min\{\mu_P(u); \mu_Q(v)\}$ for fuzzy numbers P and Q . Thus, if μ and ν are two fuzzy numbers, their minimum $\mu \wedge \nu$ and maximum $\mu \vee \nu$ are

$$(\mu \wedge \nu)(x) = \sup_{\min\{u,v\}=x} \min\{\mu(u); \nu(v)\}, \quad (\mu \vee \nu)(x) = \sup_{\max\{u,v\}=x} \min\{\mu(u); \nu(v)\}.$$

We will also use the complement operator: $(\neg\mu)(x) = \mu(1 - x)$.

A *type-2 fuzzy set* B over the universal set X is given by its membership function $\mu_B: X \times [0; 1] \mapsto [0; 1]$, which now takes two arguments. The value $\mu_B(x, u)$ may be thought of as a degree of certainty that the membership degree of x in B should be equal to u . If the element x is fixed, then $\mu_B(x, u)$ can be viewed as a fuzzy number over $[0; 1]$. Thus, it can be said that in type-2 fuzzy sets the degree of membership of every element is given by a fuzzy number, while in type-1 fuzzy sets membership degrees were ordinary real numbers. Hence, it is natural to define union and intersection of type-2 fuzzy sets via maximum and minimum of fuzzy numbers.

We will say that a type-2 fuzzy set B has *normal (convex, etc.) fuzzy grades* if for every $x \in X$ its membership function $\mu_B(x, u)$ is a normal (convex, etc.) fuzzy number over $[0; 1]$.

Following [5], we define a *meet* of type-2 fuzzy sets A and B as the type-2 fuzzy set $A \sqcap B$ given by $\mu_{A \sqcap B}(x, u) = \mu_A(x, u) \wedge \mu_B(x, u)$. Their *join* $A \sqcup B$ is $\mu_{A \sqcup B}(x, u) = \mu_A(x, u) \vee \mu_B(x, u)$. The complement $\neg A$ of a type-2 fuzzy set A is $\mu_{\neg A}(x, u) = \mu_A(x, 1-u)$. The inclusion relation is defined as follows: $A \sqsubseteq B$, if $A \sqcap B = A$ and $A \sqcup B = B$. This definition can be transferred to fuzzy numbers as well: a fuzzy number μ is said to be *not greater than* a fuzzy number ν , $\mu \sqsubseteq \nu$, if $\mu \wedge \nu = \mu$ and $\mu \vee \nu = \nu$. For normal and convex fuzzy numbers the conditions $\mu \wedge \nu = \mu$ and $\mu \vee \nu = \nu$ are equivalent [5].

3. Definitions. We will employ Zadeh's extension principle to define infimum and supremum of a set of fuzzy numbers. Let $\mu_i, i \in I$, be fuzzy numbers. The index set I may be finite or infinite, countable or uncountable. Their *infimum* $\bigwedge_{i \in I} \mu_i$ and *supremum* $\bigvee_{i \in I} \mu_i$ are fuzzy numbers with the following membership functions:

$$\left(\bigwedge_{i \in I} \mu_i \right) (x) = \sup_{\inf_{i \in I} x_i = x} \inf_{i \in I} \mu_i(x_i), \quad \left(\bigvee_{i \in I} \mu_i \right) (x) = \sup_{\sup_{i \in I} x_i = x} \inf_{i \in I} \mu_i(x_i).$$

The suprema are taken over all possible sets of x_i that have the specified infimum or supremum.

Using these operations, it is possible to define the join and meet of an arbitrary number of type-2 fuzzy sets. Let $A_i, i \in I$, be type-2 fuzzy sets. Their meet $\prod_{i \in I} A_i$ and join $\bigsqcup_{i \in I} A_i$ are type-2 fuzzy sets with the membership functions given by

$$\mu_{\prod_{i \in I} A_i}(x, u) = \bigwedge_{i \in I} \mu_{A_i}(x, u), \quad \mu_{\bigsqcup_{i \in I} A_i}(x, u) = \bigvee_{i \in I} \mu_{A_i}(x, u). \quad (1)$$

4. De Morgan's laws. Hereafter, without loss of generality, we will suppose that all fuzzy numbers under consideration are defined over $[0; 1]$.

Lemma 1. For any fuzzy numbers $\mu_i, i \in I$,

$$\neg \bigwedge_{i \in I} \mu_i = \bigvee_{i \in I} \neg \mu_i, \quad \neg \bigvee_{i \in I} \mu_i = \bigwedge_{i \in I} \neg \mu_i.$$

Proof. By definition, $\left(\neg \bigwedge_{i \in I} \mu_i \right) (x) = \left(\bigwedge_{i \in I} \mu_i \right) (1-x) = \sup_{\inf_{i \in I} x_i = 1-x} \inf_{i \in I} \mu_i(x_i) =$

$\sup_{1-\inf_{i \in I} x_i = x} \inf_{i \in I} \mu_i(x_i) = \sup_{\sup_{i \in I} (1-x_i) = x} \inf_{i \in I} \neg \mu_i(1-x_i) = \left(\bigvee_{i \in I} \neg \mu_i \right) (x)$. The second equation can be proved similarly. \square

Theorem 1. For any type-2 fuzzy sets $A_i, i \in I$,

$$\neg \prod_{i \in I} A_i = \bigsqcup_{i \in I} \neg A_i, \quad \neg \bigsqcup_{i \in I} A_i = \prod_{i \in I} \neg A_i.$$

P r o o f. From (1), $\mu_{\neg \prod_{i \in I} A_i}(x, u) = \neg \bigwedge_{i \in I} \mu_{A_i}(x, u)$, $\mu_{\neg \bigsqcup_{i \in I} A_i}(x, u) = \bigvee_{i \in I} \neg \mu_{A_i}(x, u)$, so the first equation directly follows from lemma 1. So does the second equation. \square

Thus, De Morgan's laws hold for any number of type-2 fuzzy sets.

In the following discussion, we will concentrate on properties of infimum of fuzzy numbers, and, therefore, meet of type-2 fuzzy sets. Using De Morgan's laws, it will be easy to transfer the obtained results to supremum of fuzzy numbers and join of type-2 fuzzy sets.

5. Properties of infimum of fuzzy numbers. In this section we will study whether the infimum operation preserves the properties of normality, convexity and upper semicontinuity.

Lemma 2. If $\mu_i, i \in I$, are normal fuzzy numbers, then their infimum $\bigwedge_{i \in I} \mu_i$ and supremum $\bigvee_{i \in I} \mu_i$ are also normal fuzzy numbers.

P r o o f. Denote $\mu = \bigwedge_{i \in I} \mu_i$. Take arbitrary $\varepsilon > 0$. As μ_i are normal, there exist such x_i that $\mu_i(x_i) > 1 - \varepsilon$. Then $\inf_{i \in I} \mu_i(x_i) \geq 1 - \varepsilon$. If we denote $\inf_{i \in I} x_i = x$, then $\mu(x) = \sup_{i \in I} \inf_{y_i = x} \mu_i(y_i) \geq \inf_{i \in I} \mu_i(x_i) \geq 1 - \varepsilon$. As this is true for any $\varepsilon > 0$, we may conclude that $\sup \mu(x) = 1$. The proof for supremum is similar. \square

Lemma 3. If $\mu_i, i \in I$, are strongly normal fuzzy numbers, then so are their infimum $\bigwedge_{i \in I} \mu_i$ and supremum $\bigvee_{i \in I} \mu_i$.

P r o o f. Consider the infimum $\mu = \bigwedge_{i \in I} \mu_i$, the reasoning for supremum is similar. As μ_i are strongly normal, there exist such x_i^* that $\mu_i(x_i^*) = 1$. Let $x^* = \inf_{i \in I} x_i^*$. Then $\mu(x^*) = \sup_{i \in I} \inf_{x_i = x^*} \mu_i(x_i) \geq \inf_{i \in I} \mu_i(x_i^*) = 1$. \square

Lemma 4. If $\mu_i, i \in I$, are convex fuzzy numbers, then their infimum $\bigwedge_{i \in I} \mu_i$ and supremum $\bigvee_{i \in I} \mu_i$ are also convex fuzzy numbers.

P r o o f. Consider the infimum $\mu = \bigwedge_{i \in I} \mu_i$, the proof for supremum is similar.

A fuzzy number μ_i over $[0; 1]$ is convex if and only if its membership function is nondecreasing on $[0; x_i^*]$ and nonincreasing on $[x_i^*; 1]$ or nondecreasing on $[0; x_i^*]$ and nonincreasing on $(x_i^*; 1]$ for some x_i^* . Let $x^* = \inf_{i \in I} x_i^*$.

Take $x < y < x^*$ and suppose that $\mu(x) > \mu(y)$. Let $\varepsilon = \frac{1}{2}(\mu(x) - \mu(y)) > 0$. As $\mu(x) = \sup_{i \in I} \inf_{x_i = x} \mu_i(x_i)$, there exist such x_i that $\inf_{i \in I} x_i = x$ and $\inf_{i \in I} \mu_i(x_i) > \mu(x) - \varepsilon$.

Let $y_i = \max\{x_i; y\} \geq x_i$. Since $x^* \leq x_i^*$, all functions μ_i are nondecreasing on $[0; x^*]$. Hence, $\mu_i(y_i) \geq \mu_i(x_i)$, and $\inf_{i \in I} \mu_i(y_i) \geq \inf_{i \in I} \mu_i(x_i)$. As $\inf_{i \in I} \mu_i(x_i) = x < y$, there exists such index j that $x_j < y$, and then $y_j = y$. At the same time, $y_i \geq y$ for all $i \in I$. Thus, $\inf_{i \in I} y_i = y$. Therefore, $\mu(y) \geq \inf_{i \in I} \mu_i(y_i) \geq \inf_{i \in I} \mu_i(x_i) > \mu(x) - \varepsilon = \frac{1}{2}(\mu(x) + \mu(y)) > \mu(y)$. This contradiction proves that μ is nondecreasing on $[0; x^*]$.

Consider the set $J(x) = \{i \in I: x_i^* < x\}$. All functions μ_j with $j \in J(x)$ are nonincreasing on $[x; 1]$. If $x > x^* = \inf_{i \in I} x_i^*$, then $J(x) \neq \emptyset$. Take $x^* < x < y$ and suppose that $\mu(x) < \mu(y)$. Let $\varepsilon = \frac{1}{2}(\mu(y) - \mu(x)) > 0$. There exist such y_i that $\inf_{i \in I} y_i = y$ and $\inf_{i \in I} \mu_i(y_i) > \mu(y) - \varepsilon$. Let $x_i = x$ for $i \in J(x)$, and $x_i = y_i$ for $i \notin J(x)$. As $J(x) \neq \emptyset$, $\inf_{i \in I} x_i = x$. Then $\mu(x) \geq \inf_{i \in I} \mu_i(x_i)$. For $i \in J(x)$, as $y_i \geq \inf_{i \in I} y_i = y > x = x_i$, due to μ_i being nonincreasing on $[x; 1]$, $\mu_i(x_i) \geq \mu_i(y_i)$. For $i \notin J(x)$, we have $x_i = y_i$, so $\mu_i(x_i) = \mu_i(y_i)$. Thus, $\inf_{i \in I} \mu_i(x_i) \geq \inf_{i \in I} \mu_i(y_i)$. Collecting all inequalities, we obtain $\mu(x) > \mu(y) - \varepsilon = \mu(x) + \varepsilon$, a contradiction. Hence, the function μ must be nonincreasing on $(x^*; 1]$.

Let $K = \{i \in I: x_i^* = x^*\}$. If $i \notin K$, then $x_i^* > x^*$, and the function μ_i is nondecreasing on $[0; x^*]$.

Consider the case where for $\forall i \in K$ the functions μ_i are nondecreasing on $[0; x^*]$, or $K = \emptyset$. Then all functions μ_i , $i \in I$, are nondecreasing on $[0; x^*]$. Take $x < x^*$ and suppose that $\mu(x) > \mu(x^*)$. Let $\varepsilon = \frac{1}{2}(\mu(x^*) - \mu(x)) > 0$. Then there exist x_i such that $\inf_{i \in I} x_i = x$ and $\inf_{i \in I} \mu_i(x_i) > \mu(x) - \varepsilon$. As $x < x^*$, there will be indices $i \in I$ for which $x_i < x^*$. For these i let $y_i = x^*$. Then $\mu_i(y_i) \geq \mu_i(x_i)$. For all other indices, in other words, when $x_i \geq x^*$, let $y_i = x_i$. Then $\inf_{i \in I} y_i = x^*$, and $\mu(x^*) \geq \inf_{i \in I} \mu_i(y_i) \geq \inf_{i \in I} \mu_i(x_i) > \mu(x) - \varepsilon = \mu(x^*) + \varepsilon$, a contradiction. Thus, in this case the function μ is nondecreasing on $[0; x^*]$.

Now consider the case where for some $k \in K$ the function μ_k is nondecreasing on $[0; x^*)$ and nonincreasing on $[x^*; 1]$. Suppose that for some $y > x^*$ we have $\mu(x^*) < \mu(y)$. Let $\varepsilon = \frac{1}{2}(\mu(y) - \mu(x^*)) > 0$. There exist such y_i that $\inf_{i \in I} y_i = y$ and $\inf_{i \in I} \mu_i(y_i) > \mu(y) - \varepsilon$. Let $x_i = y_i$ for $i \neq k$, and $x_k = x^* < y \leq y_k$, so that $\mu_k(x_k) \geq \mu_k(y_k)$. Then $\inf_{i \in I} x_i = x^*$, and $\mu(x^*) \geq \inf_{i \in I} \mu_i(x_i) \geq \inf_{i \in I} \mu_i(y_i) > \mu(y) - \varepsilon > \mu(x) + \varepsilon$, a contradiction. Thus, in this case the function μ is nonincreasing on $[x^*; 1]$.

Summing up, the function μ is either nondecreasing on $[0; x^*]$ and nonincreasing on $(x^*; 1]$, or nondecreasing on $[0; x^*)$ and nonincreasing on $[x^*; 1]$. Thus, μ is convex. \square

Theorem 2. *The join and meet of any number of type-2 fuzzy sets with normal (strongly normal, convex) fuzzy grades are also type-2 fuzzy sets with normal (strongly normal, convex) fuzzy grades.*

Proof follows from (1) and lemmas 2–4. \square

Unfortunately, the set of upper semicontinuous fuzzy numbers is not closed with respect to the infimum and supremum operators, as the following example demonstrates. Let

$$\mu_i(x) = \begin{cases} 1, & 0 \leq x \leq \frac{1}{2} - \frac{1}{i}, \\ 0, & \frac{1}{2} - \frac{1}{i} < x < \frac{2}{3}, \\ 1, & \frac{2}{3} \leq x \leq 1, \end{cases}$$

for $i \in \mathbb{N}$. It is easy to verify that

$$\left(\bigwedge_{i=1}^{\infty} \mu_i \right) (x) = \begin{cases} 1, & 0 \leq x < \frac{1}{2}, \\ 0, & \frac{1}{2} \leq x < \frac{2}{3}, \\ 1, & \frac{2}{3} \leq x \leq 1. \end{cases}$$

Thus, despite every μ_i being upper semicontinuous, their infimum is not upper semicontinuous.

Lemma 5. If $\mu_i, i \in I$, are normal convex upper semicontinuous fuzzy numbers, then their infimum $\bigwedge_{i \in I} \mu_i$ and supremum $\bigvee_{i \in I} \mu_i$ are strongly normal, convex, and upper semicontinuous.

Proof. Consider the infimum $\mu = \bigwedge_{i \in I} \mu_i$, as the proof for supremum is similar.

As normal upper semicontinuous fuzzy numbers are strongly normal, there exist x_i^* such that $\mu_i(x_i^*) = 1$. Let $x^* = \inf_{i \in I} x_i^*$. Then $\mu(x^*) = 1$.

From previous lemmas it follows that μ is strongly normal and convex, so it remains to show that μ is upper semicontinuous. Consider an α -cut $A = \{x: \mu(x) \geq \alpha\}$. As μ is strongly normal, it is nonempty. Let $x' = \inf_{\mu(x) \geq \alpha} x$, and $x'' = \sup_{\mu(x) \geq \alpha} x$. If $x' = x''$, then the set $A = \{x'\}$ consists of a single point and is thus closed. Consider the case $x' < x''$. By convexity of μ , $(x'; x'') \subseteq A$. As $\mu(x^*) = 1 \geq \alpha$, $x' \leq x^* \leq x''$.

For every $i \in I$ we define $B_i = \{x: \mu_i(x) \geq \alpha\}$. Since all μ_i are upper semicontinuous, these sets are closed, and since all μ_i are strongly normal and convex, $B_i = [a_i; b_i]$, and $a_i \leq x_i^* \leq b_i$.

If $\exists j \in I: \mu_j(x') \geq \alpha$, then, taking $x_j = x'$ and $x_i = x_i^* \geq x^* \geq x'$ for $i \neq j$, we obtain $\inf_{i \in I} x_i = x'$ and $\inf_{i \in I} \mu_i(x_i) = \mu_j(x') \geq \alpha$, so $\mu(x') \geq \alpha$. Suppose that $\mu_j(x') < \alpha$ for all $j \in I$. As $x' \leq x^* \leq x_i^* \leq b_i$, $x' < a_i$. Consider $a = \inf_{i \in I} a_i$. We have $\mu_i(a_i) \geq \alpha$, so $\inf_{i \in I} \mu_i(a_i) \geq \alpha$. Since $x' < a_i$ for $\forall i \in I$, $x' \leq a$. If $a = x'$, then immediately $\mu(x') \geq \alpha$.

Consider the case $x' < a$. Take $\varepsilon > 0$, $y_n = x' + \frac{1}{n}$ for $n > n_0 = \max\left\{\frac{1}{a-x'}, \frac{1}{x''-x'}\right\}$, so that $x' < y_n < \min\{a; x''\}$. As $y_{2n} \in (x'; x'')$, $\mu(y_{2n}) \geq \alpha$. Then there exist x_i^n such that $\inf_{i \in I} x_i^n = y_{2n}$ and $\inf_{i \in I} \mu_i(x_i^n) > \alpha - \varepsilon$. Since $y_{2n} < y_n$, there exists such $j_n \in I$ that $y_{2n} \leq x_{j_n}^n < y_n$ and $\mu_{j_n}(x_{j_n}^n) > \alpha - \varepsilon$. Let now $x_i = \inf_{j_n=i} x_{j_n}^n$ for $i \in \bigcup_{n > n_0} \{j_n\}$, and $x_i = x_i^*$ for the remaining indices $i \in I \setminus \bigcup_{n > n_0} \{j_n\}$. If $i \in \bigcup_{n > n_0} \{j_n\}$, then $\mu_i(x_{j_n}^n) > \alpha - \varepsilon$ whenever $j_n = i$, so using upper semicontinuity we get $\mu_i(x_i) \geq \alpha - \varepsilon$. For the remaining indices $\mu_i(x_i) = 1$. Thus, $\inf_{i \in I} \mu_i(x_i) \geq \alpha - \varepsilon$, while $\inf_{i \in I} x_i = \inf_{n > n_0} x_{j_n}^n = x'$. Therefore, $\mu(x') \geq \alpha$.

Consider the set $J = \{i \in I: x_i^* < x''\}$. If $J = \emptyset$, $x'' \leq x_i^*$ for all $i \in I$, hence, $x'' \leq x^*$. But $x^* \leq x''$, so $x^* = x''$, and $\mu(x'') = \mu(x^*) = 1 \geq \alpha$. Consider the case $J \neq \emptyset$. Let $b = \inf_{j \in J} b_j$. If $b \geq x''$, then $b_j \geq x'' > x_j^*$ for every $j \in J$, so $\mu_j(x'') \geq \mu_j(b_j) \geq \alpha$ as μ_j are nonincreasing on $(x_j^*; 1]$. Then, taking $x_i = x''$ for $i \in J$ and $x_i = x_i^* \geq x''$ for $i \notin J$, we obtain $\inf_{i \in I} \mu_i(x_i) \geq \alpha$ and $\inf_{i \in I} x_i = x''$, so $\mu(x'') \geq \alpha$. Consider the case $b < x''$. Take $y \in (\max\{x'; b\}; x'')$. As $b < y$, $\exists j \in J: b_j < y$. Then $\mu_j(y) < \alpha$. As μ_j is nonincreasing on $(x_j^*; 1]$, $\mu_j(x_j) \leq \mu_j(y)$ for $\forall x_j \geq y$. Then for any x_i such that $\inf_{i \in I} x_i = y$ we will have $x_j \geq y$, so $\inf_{i \in I} \mu_i(x_i) \leq \mu_j(x_j) \leq \mu_j(y)$. Then $\mu(y) \leq \mu_j(y) < \alpha$. But $y \in (x'; x'')$, so $\mu(y) \geq \alpha$, a contradiction. Thus, $b < x''$ is impossible.

Summing up, we have shown that $\mu(x') \geq \alpha$ and $\mu(x'') \geq \alpha$, so $A = [x'; x'']$ is a closed set. Thus, μ is upper semicontinuous. \square

Theorem 3. The join and meet of type-2 fuzzy sets with normal convex upper semicontinuous fuzzy grades are type-2 fuzzy sets with strongly normal convex upper semicontinuous fuzzy grades.

Proof follows from (1) and lemma 5. \square

6. Idempotency of infimum. Consider the following example. Let

$$\mu(x) = \begin{cases} 0, & 0 \leq x \leq \frac{1}{3}, \\ 1, & \frac{1}{3} < x < \frac{2}{3}, \\ 0, & \frac{2}{3} \leq x \leq 1. \end{cases}$$

It is a strongly normal and convex fuzzy number, but it is not upper semicontinuous. It is easy to verify that $\mu \wedge \mu = \mu$. But if we take an infinite number of identical fuzzy numbers $\mu_i = \mu$, $i \in N$, and compute their infimum $\nu = \bigwedge_{i=1}^{\infty} \mu_i$, then we will find that $\bigwedge_{i=1}^{\infty} \mu_i \neq \mu$:

$$\nu(x) = \begin{cases} 0, & 0 \leq x < \frac{1}{3}, \\ 1, & \frac{1}{3} \leq x < \frac{2}{3}, \\ 0, & \frac{2}{3} \leq x \leq 1. \end{cases}$$

Indeed, due to infinite number of arguments, we may take $x_i = \frac{1}{3} + \frac{1}{i}$, so that $\inf_{i \in N} x_i = \frac{1}{3}$, $\inf_{i \in N} \mu_i(x_i) = 1$, and thus yield $\nu(\frac{1}{3}) = 1$. The infimum ν is still not upper semicontinuous, however. This example demonstrates that using not upper semicontinuous fuzzy numbers may violate one of intuitive properties of infimum, namely, idempotency: for real numbers, if $x_i = x$ for $\forall i \in N$, then $\inf_{i \in N} x_i = x$, but this may not hold for fuzzy numbers unless they are upper semicontinuous.

Let $J = N$, and $I_j = \{1\}$ for every $j \in J$. Observe that $\bigcup_{j \in J} I_j = \{1\}$ is a finite set. Let $\mu_1 = \mu$ from the previous example. Then we have $\bigwedge_{j \in J} \bigwedge_{i \in I_j} \mu_i \neq \bigwedge_{i \in \bigcup_{j \in J} I_j} \mu_i$. Thus, even such

seemingly obvious properties as independence of infimum from reordering or grouping its arguments must be carefully examined.

Lemma 6. Let μ_i , $i \in I$, be fuzzy numbers, and $I = \bigcup_{j \in J} I_j$. Then if J is a finite set, or all the fuzzy numbers μ_i are upper semicontinuous, $\bigwedge_{j \in J} \bigwedge_{i \in I_j} \mu_i = \bigwedge_{i \in I} \mu_i$, and $\bigvee_{j \in J} \bigvee_{i \in I_j} \mu_i = \bigvee_{i \in I} \mu_i$.

Proof. Denote $\nu_j = \bigwedge_{i \in I_j} \mu_i$, $\nu = \bigwedge_{j \in J} \nu_j$, $\mu = \bigwedge_{i \in I} \mu_i$.

Suppose first that $\mu(x) > \nu(x)$. Take $\varepsilon = \frac{1}{2}(\mu(x) - \nu(x)) > 0$. Then there exist x_i such that $\inf_{i \in I} x_i = x$, $\inf_{i \in I} \mu_i(x_i) > \mu(x) - \varepsilon$. For $y_j = \inf_{i \in I_j} x_i$ we will have $\nu_j(y_j) \geq \inf_{i \in I_j} \mu_i(x_i)$, so $\inf_{j \in J} \nu_j(y_j) \geq \inf_{i \in I} \mu_i(x_i) > \mu(x) - \varepsilon$. As for any $\delta > 0$ there exists $i \in I$ for which $x_i < x + \delta$, there exists $j \in J$ such that the index $i \in I_j$, so $y_j \leq x_i < x + \delta$. Thus, $\inf_{j \in J} y_j = x$. Therefore, $\nu(x) \geq \inf_{j \in J} \nu_j(y_j) > \mu(x) - \varepsilon = \nu(x) + \varepsilon$, a contradiction.

Suppose now that $\mu(x) < \nu(x)$. Take $\varepsilon = \frac{1}{4}(\nu(x) - \mu(x)) > 0$. There exist such x_j that $\inf_{j \in J} x_j = x$ and $\inf_{j \in J} \nu_j(x_j) > \nu(x) - \varepsilon$. Then, for every $j \in J$, there exist such x_{ji} that $\inf_{i \in I_j} x_{ji} = x_j$ and $\inf_{i \in I_j} \mu_i(x_{ji}) > \nu_j(x_j) - \varepsilon > \nu(x) - 2\varepsilon$. Let $y_i = \inf_{j: i \in I_j} x_{ji}$. Since $x_{ji} \geq x_j \geq x$, $y_i \geq x$. For any $\delta > 0$ we can find such index j that $x_j < x + \delta$, and then such $i \in I_j$ that $x_{ji} < x_j + \delta < x + 2\delta$. Then $y_i \leq x_{ji} < x + 2\delta$. Thus, $\inf_{i \in I} y_i = x$. If μ_i are upper semicontinuous, then for every $i \in I$, for all j such that $i \in I_j$, from

$\mu_i(x_{ji}) \geq \inf_{i \in I_j} \mu_i(x_{ji}) \geq \nu(x) - 2\varepsilon$ we may conclude that $\mu_i(y_i) \geq \nu(x) - 2\varepsilon$. If J is finite, $y_i = x_{ji}$ for some $j \in J$, so again $\mu_i(y_i) = \mu_i(x_{ji}) \geq \inf_{i \in I_j} \mu_i(x_{ji}) \geq \nu(x) - 2\varepsilon$. Then $\mu(x) \geq \inf_{i \in I} \mu_i(y_i) \geq \nu(x) - 2\varepsilon = \mu(x) + 2\varepsilon$, a contradiction.

Thus, $\mu = \nu$, as desired. The second formula can be proved in a similar way. \square

Theorem 4. Let $A_i, i \in I$, be type-2 fuzzy sets, and $I = \bigcup_{j \in J} I_j$. If J is finite, or the type-2 fuzzy sets A_i have upper semicontinuous fuzzy grades, then $\bigcup_{j \in J} \bigcup_{i \in I_j} A_i = \bigcup_{i \in I} A_i$,

$$\prod_{j \in J} \prod_{i \in I_j} A_i = \prod_{i \in I} A_i.$$

Proof follows from (1) and lemma 6. \square

Theorem 5. Let $A_i, i \in I$, be type-2 fuzzy sets with normal convex fuzzy grades. Then

- 1) $\prod_{i \in I} A_i \subseteq A_j \subseteq \bigcup_{i \in I} A_i$ for $\forall j \in I$;
- 2) $\prod_{i \in I} \mu_i \subseteq \prod_{i \in J} \mu_i$ for any $J \subseteq I$;
- 3) $\bigcup_{i \in J} \mu_i \subseteq \bigcup_{i \in I} \mu_i$ for any $J \subseteq I$.

Proof. By theorem 4, $A_j \cap \prod_{i \in I} A_i = \prod_{i \in \{j\} \cup I} A_i = \prod_{i \in I} A_i$, and due to normality and convexity of fuzzy grades $\prod_{i \in I} A_i \subseteq A_j$. The other properties can be proved similarly. \square

Theorem 6. For type-2 fuzzy sets with normal convex upper semicontinuous fuzzy grades:

- 1) if $A \subseteq B_i$ for $\forall i \in I$, then $A \subseteq \prod_{i \in I} B_i$;
- 2) if $B_i \subseteq A$ for $\forall i \in I$, then $\bigcup_{i \in I} B_i \subseteq A$;
- 3) if $A_i \subseteq B_i$ for $\forall i \in I$, then $\prod_{i \in I} A_i \subseteq \prod_{i \in I} B_i$, and $\bigcup_{i \in I} A_i \subseteq \bigcup_{i \in I} B_i$.

Proof. Denote $A = B_0$. If $A \subseteq B_i$, then $B_0 \cap B_i = B_0$. Using theorem 4, we obtain $A \cap \prod_{i \in I} B_i = \prod_{i \in I \cup \{0\}} B_i = \prod_{i \in \bigcup_{j \in I} \{0; j\}} B_i = \prod_{j \in I} (B_0 \cap B_j) = \prod_{j \in I} B_0 = \prod_{j \in I} \prod_{i \in \{0\}} B_i =$

$\prod_{i \in \bigcup_{j \in I} \{0\}} B_i = B_0 = A$. The second property is similar. Finally, by theorem 5, $\prod_{i \in I} A_i \subseteq$

$A_j \subseteq B_j \subseteq \bigcup_{i \in I} B_i$ for any $j \in I$. Then by previous properties we get the remaining inclusions. \square

7. Formulae for join and meet of type-2 fuzzy sets. Let $\mu_i, i \in I = \{1, 2, \dots, n\}$, be strongly normal convex fuzzy numbers. Let x_i^* be such points that $\mu_i(x_i^*) = 1, i \in I$. Without loss of generality, suppose that $x_1^* \leq x_2^* \leq \dots \leq x_n^*$. Then infimum and supremum of these numbers can be computed as follows [7]:

$$\left(\bigwedge_{i=1}^n \mu_i \right) (x) = \begin{cases} \max_{i=1, \dots, n} \mu_i(x), & x < x_1^*, \\ \min_{i=1, \dots, k} \mu_i(x), & x_k^* \leq x < x_{k+1}^*, \\ \min_{i=1, \dots, n} \mu_i(x), & x \geq x_n^*, \end{cases}$$

$$\left(\bigvee_{i=1}^n \mu_i\right)(x) = \begin{cases} \min_{i=1, \dots, n} \mu_i(x), & x < x_1^*, \\ \min_{i=k+1, \dots, n} \mu_i(x), & x_k^* \leq x < x_{k+1}^*, \\ \max_{i=1, \dots, n} \mu_i(x), & x \geq x_n^*. \end{cases}$$

These formulae at the same time allow to compute join and meet of a finite number of type-2 fuzzy sets, as these operations by definition are reduced to maximum and minimum of fuzzy numbers corresponding to each element of the universal set.

Consider the following example. Let $\mu_n(x) = 1 - (1 - \sqrt[n]{x})^n$, $n \in N$, and $\mu = \bigwedge_{n=1}^{\infty} \mu_n$. Take arbitrary $\varepsilon > 0$. If $x_n = 2^{-n}$, then $\lim_{n \rightarrow \infty} \mu_n(x_n) = \lim_{n \rightarrow \infty} (1 - 2^{-n}) = 1$. Then for some $n_0 \in N$ we will have $\mu_n(x_n) > 1 - \varepsilon$ for $\forall n \geq n_0$. Let $x_i = 2^{-i}$ for $i \geq n_0$, and $x_i = 1$ for $i < n_0$. Then $\inf_{i \in N} x_i = 0$, and $\inf_{i \in N} \mu_i(x_i) \geq 1 - \varepsilon$, so $\mu(0) \geq 1 - \varepsilon$. Due to arbitrariness of ε , $\mu(0) = 1$. Now note that $\mu_n(0) = 0$ for $\forall n \in N$, so $\sup_{n \in N} \mu_n(0) \neq \mu(0)$. Thus, the given formulae cannot be naively generalized to the case of infinite number of fuzzy numbers.

Lemma 7. Let μ_i , $i \in I$, be normal convex upper semicontinuous fuzzy numbers, and $\mu = \bigwedge_{i \in I} \mu_i$. Then

$$\mu(x) = \begin{cases} \overline{\lim}_{y \rightarrow x+0} \sup_{i \in I} \mu_i(y), & \forall i \in I \nexists y \leq x: \mu_i(y) = 1, \\ \inf_{i \in I: \exists y \leq x: \mu_i(y)=1} \mu_i(x), & \exists i \in I \exists y \leq x: \mu_i(y) = 1. \end{cases} \quad (2)$$

Proof. As normal upper semicontinuous fuzzy numbers are strongly normal, there exist x_i such that $\mu_i(x_i) = 1$. Let $x_i^* = \inf_{\mu_i(x_i)=1} x_i$. By upper semicontinuity, $\mu_i(x_i^*) = 1$. Let also $x^* = \inf_{i \in I} x_i^*$.

Consider $x < x^*$. By choice of x^* we have $\forall i \in I \nexists y \leq x: \mu_i(y) = 1$. Since all μ_i are convex, they are nondecreasing on $[x; x^*]$. Then so is $\varphi(y) = \sup_{i \in I} \mu_i(y)$. Indeed, if for some x', x'' such that $x \leq x' < x'' \leq x^*$ we had $\varphi(x') = \sup_{i \in I} \mu_i(x') > \sup_{i \in I} \mu_i(x'') = \varphi(x'')$, then for $\varepsilon = \frac{1}{2}(\varphi(x') - \varphi(x'')) > 0$ there would exist such $j \in I$ that $\mu_j(x') > \varphi(x') - \varepsilon = \varphi(x'') + \varepsilon > \varphi(x'') \geq \mu_j(x'')$, so μ_j would not be nondecreasing. Furthermore, $\varphi(y) \geq 0$, so the function $\varphi(y)$ is monotone and bounded on $[x; x^*]$. Therefore, $\psi(x) = \lim_{y \rightarrow x+0} \varphi(y)$ exists. As $\varphi(y)$ is nondecreasing, $\psi(x) \leq \varphi(y)$ for $\forall y \in [x; x^*]$.

Let $y_n = x + \frac{x^* - x}{2^n}$, $n \in N$, be a monotonically decreasing sequence, $y_n \rightarrow x + 0$ as $n \rightarrow \infty$. Since $y_n \in (x; x^*)$, $\varphi(y_n) \geq \psi(x)$. Take arbitrary $\varepsilon > 0$. Then for every $n \in N$ we can find $i_n \in I$ such that $\mu_{i_n}(y_n) > \varphi(y_n) - \varepsilon \geq \psi(x) - \varepsilon$. Denote $I_\varepsilon = \{i_n: n \in N\}$. Let $x_i = \inf_{n \in N: i_n=i} y_n$ for $i \in I_\varepsilon$, and $x_i = x_i^*$ for $i \notin I_\varepsilon$. Since $y_n \rightarrow x + 0$, $\inf_{i \in I} x_i = x$. For $i \in I_\varepsilon$ we have $\mu_i(y_n) \geq \psi(x) - \varepsilon$ for $n \in N: i_n = i$, so due to μ_i being upper semicontinuous, $\mu_i(x_i) \geq \psi(x) - \varepsilon$. For other indices $i \notin I_\varepsilon$, simply $\mu_i(x_i) = 1$. Thus, $\mu(x) \geq \inf_{i \in I} \mu_i(x_i) \geq \psi(x) - \varepsilon$. Due to arbitrariness of ε , $\mu(x) \geq \psi(x)$.

Suppose that $\mu(x) > \psi(x)$. Let $\varepsilon = \frac{1}{2}(\mu(x) - \psi(x)) > 0$. Then $\exists \delta > 0: \forall y \in (x; x + \delta) \Rightarrow |\varphi(y) - \varphi(x)| < \varepsilon$. Hence, $\varphi(y) < \psi(x) + \varepsilon = \mu(x) - \varepsilon$ for $\forall y \in (x; x + \delta)$. As φ is nondecreasing, $\varphi(x) \leq \varphi(y) < \mu(x) - \varepsilon$. Since $\mu(x) = \sup_{i \in I} \inf_{x_i=x} \mu_i(x_i)$, there exist

such x_i that $\inf_{i \in I} x_i = x$ and $\inf_{i \in I} \mu_i(x_i) > \mu(x) - \varepsilon$. For the found δ then $\exists j: x_j < x + \delta$, so $\varphi(x_j) < \mu(x) - \varepsilon < \inf_{i \in I} \mu_i(x_i) \leq \mu_j(x_j) \leq \sup_{i \in I} \mu_i(x_j) = \varphi(x_j)$, a contradiction. Thus, $\mu(x) = \psi(x)$.

Consider now $x = x^*$. Taking $x_i = x_i^*$, we get $\inf_{i \in I} x_i = x^*$ and $\inf_{i \in I} \mu_i(x_i) = 1$, so $\mu(x^*) = 1$.

If $\exists i \in I: x_i^* = x^*$, then it is the case $\exists i \in I, \exists y \leq x: \mu_i(y) = 1$. For $i \in I$, if $\exists y \leq x = x^*: \mu_i(y) = 1$, then $x^* \leq x_i^* \leq y$, so $y = x^*$, so $\mu_i(x^*) = 1$. Then $\inf_{i \in I: \exists y \leq x: \mu_i(y)=1} \mu_i(x) = 1 = \mu(x)$.

If $x_i^* > x^*$ for $\forall i \in I$, then, obviously, $\forall i \in I \nexists y \leq x: \mu_i(y) = 1$. Since $x^* = \inf_{i \in I} x_i^*$, we may choose a sequence $x_n^* \rightarrow x^* + 0$. Then $\sup_{i \in I} \mu_i(x_n^*) \geq \mu_n(x_n^*) = 1$, so $\overline{\lim}_{y \rightarrow x^* + 0} \sup_{i \in I} \mu_i(y) = 1 = \mu(x^*)$.

Finally, consider $x > x^*$. Let $J(x) = \{i \in I: x_i^* \leq x\}$. As $x^* = \inf_{i \in I} x_i^* \exists i \in I: x_i^* < x$, so $J(x) \neq \emptyset$. Besides, the condition $x_i^* \leq x$ is equivalent to $\exists y \leq x: \mu_i(y) = 1$, so the infimum in (2) is taken over $J(x)$ exactly.

Let $x_i = x$ for $i \in J(x)$, and $x_i = x_i^*$ for $i \notin J(x)$. Then $\inf_{i \in I} x_i = x$ and $\inf_{i \in I} \mu_i(x_i) = \inf_{i \in J(x)} \mu_i(x)$, so $\mu(x) \geq \inf_{i \in J(x)} \mu_i(x)$.

Suppose that $\mu(x) > \chi(x) = \inf_{i \in J(x)} \mu_i(x)$. Let $\varepsilon = \frac{1}{2}(\mu(x) - \chi(x)) > 0$. Then there exist such x_i that $\inf_{i \in I} x_i = x$ and $\inf_{i \in I} \mu_i(x_i) > \mu(x) - \varepsilon$. Hence, $\mu_i(x_i) > \mu(x) - \varepsilon$ for all $i \in I$. On the other hand, as $\chi(x) = \inf_{i \in J(x)} \mu_i(x) \exists j \in J(x): \mu_j(x) < \chi(x) + \varepsilon = \mu(x) - \varepsilon$. As $j \in J(x)$, $x_j^* \leq x$, so by convexity the function μ_j is nonincreasing on $[x; 1]$. As $x_j \geq x$, $\mu_j(x_j) \leq \mu_j(x) < \mu(x) - \varepsilon$, a contradiction. Thus, $\mu(x) = \chi(x)$. \square

It must be noted that we cannot completely get rid of the upper limit sign in (2), as the following example demonstrates. Let

$$\mu_n(x) = \begin{cases} 1, & x = \frac{1}{n}, \\ 0, & x \neq \frac{1}{n}, \end{cases}$$

be a fuzzy number that represents the crisp number $\frac{1}{n}$. Obviously,

$$\left(\bigwedge_{n=1}^{\infty} \mu_n \right) (x) = \begin{cases} 1, & x = 0, \\ 0, & x \neq 0, \end{cases}$$

is the fuzzy counterpart of the crisp zero. But

$$\sup_{n \in N} \mu_n(x) = \begin{cases} 1, & \frac{1}{x} \in N, \\ 0, & \frac{1}{x} \notin N, \end{cases}$$

so the ordinary limit $\lim_{x \rightarrow +0} \sup_{n \in N} \mu_n(x)$ does not exist. Fortunately, this can happen only when the membership function of the infimum equals 1. In other cases ordinary limit exists, as follows from the proof.

The presented formula (2) can be used to compute fuzzy grades for join and meet of any number of type-2 fuzzy sets, as in the following result.

Theorem 7. Let $A_i, i \in I$, be type-2 fuzzy sets with normal convex upper semicontinuous fuzzy grades. Then for any $x \in X, u \in [0; 1]$,

$$\mu_{\prod_{i \in I} A_i}(x, u) = \begin{cases} \lim_{v \rightarrow u+0} \sup_{i \in I} \mu_{A_i}(x, u), & u < u^*(x), \\ 1, & u = u^*(x), \\ \inf_{i \in J(x, u)} \mu_{A_i}(x, u), & u > u^*(x), \end{cases}$$

$$\mu_{\sqcup_{i \in I} A_i}(x, u) = \begin{cases} \inf_{i \in J'(x, u)} \mu_{A_i}(x, u), & u < u^0(x), \\ 1, & u = u^0(x), \\ \lim_{v \rightarrow u-0} \sup_{i \in I} \mu_{A_i}(x, u), & u > u^0(x), \end{cases}$$

where $u^*(x) = \inf_{i \in I} u_i^*(x), u_i^*(x) = \inf_{\mu_{A_i}(x, u)=1} u, u^0(x) = \sup_{i \in I} u_i^0(x), u_i^0(x) = \sup_{\mu_{A_i}(x, u)=1} u,$
 $J(x, u) = \{i \in I: u \geq u_i^*(x)\}, J'(x, u) = \{i \in I: u \leq u_i^0(x)\}.$

Proof. The first part directly follows from the proof of lemma 7 and (1). The second formula can be derived from the first with the use of De Morgan's laws. \square

8. Conclusions. Zadeh's extension principle gives a natural way to extend known operations on real numbers to fuzzy numbers. However, they might not retain all usual properties. We have studied an extension of infimum and supremum to fuzzy numbers. We have shown that these operations preserve normality and convexity, but do not preserve upper semicontinuity. At the same time, if fuzzy numbers are not upper semicontinuous, their infimum and supremum might behave counter-intuitively, for example, they might be not idempotent. The conditions under which familiar properties of infimum and supremum hold have been given. We have also derived formulae for computing infimum and supremum of arbitrary number of fuzzy numbers. All obtained results are applicable to type-2 fuzzy sets, as set-theoretic operations on them are defined using infimum and supremum of fuzzy numbers.

References

1. Zadeh L. A. Fuzzy sets. *Information and Control*, 1965, vol. 8, iss. 3, pp. 338–353.
2. Zadeh L. A. The concept of a linguistic variable and its application to approximate reasoning. *Information Sciences*, 1975, vol. 8, iss. 3, pp. 199–249.
3. Liang Q., Mendel J. M. MPEG VBR video traffic modeling and classification using fuzzy techniques. *IEEE Transactions on Fuzzy Systems*, 2001, vol. 9, iss. 1, pp. 183–193.
4. Hisdal E. The IF THEN ELSE statement and interval-valued fuzzy sets of higher type. *International Journal of Man-Machine Studies*, 1981, vol. 15, pp. 385–455.
5. Mizumoto M., Tanaka K. Some properties of fuzzy sets of type-2. *Fuzzy Sets and Systems*, 1976, vol. 31, pp. 312–340.
6. Noghin V. D. *Pareto set reduction. An axiomatic approach.* Cham, Springer Publ., 2018, 232 p.
7. Karnik N. N., Mendel J. M. Operations on type-2 fuzzy sets. *Fuzzy Sets and Systems*, 2001, vol. 122, pp. 327–348.

Received: December 02, 2020.

Accepted: April 05, 2021.

Author's information:

Oleg V. Baskov — PhD in Physics and Mathematics, Associate Professor; o.baskov@spbu.ru

Соединение и слияние бесконечного набора нечетких множеств типа 2*

О. В. Басков

Санкт-Петербургский государственный университет, Российская Федерация,
199034, Санкт-Петербург, Университетская наб., 7–9

Для цитирования: *Baskov O. V. Join and meet of infinite number of type-2 fuzzy sets // Вестник Санкт-Петербургского университета. Прикладная математика. Информатика. Процессы управления. 2021. Т. 17. Вып. 2. С. 108–119. <https://doi.org/10.21638/11701/spbu10.2021.201>*

Изучаются свойства соединения и слияния бесконечного количества нечетких множеств типа 2. Также исследуются тесно связанные с ними операции инфимума и супремума над нечеткими числами. Показано, что классы нечетких множеств типа 2 с нормальными или выпуклыми степенями принадлежности замкнуты относительно соединения и слияния, однако полунепрерывность сверху при этом может не сохраняться. Установлены условия, при которых справедливы такие основные свойства соединения и слияния как идемпотентность, коммутативность и ассоциативность. Доказана корректность применения инфимума и супремума к обеим частям неравенств нормальных выпуклых полунепрерывных сверху нечетких чисел. Наконец, приведена формула для нахождения функций принадлежности соединения и слияния бесконечного количества нечетких множеств типа 2.

Ключевые слова: нечеткие множества типа 2, соединение, слияние, нечеткие числа, инфимум, супремум.

Контактная информация:

Басков Олег Владимирович — канд. физ.-мат. наук, доц.; o.baskov@spbu.ru

* Работа выполнена при финансовой поддержке Российского фонда фундаментальных исследований (грант № 20-07-00298-а).