

Санкт-Петербургский государственный университет

ТАРАСОВА Мария Сергеевна

Выпускная квалификационная работа

Социологическая модель с нестандартной нормировкой

Уровень образования:

Направление 01.04.01 "Математика"

Основная образовательная программа ВМ.5832.2020 "Современная математика"

Научный руководитель:

д.ф.-м.н. профессор

Пилюгин Сергей Юрьевич

Рецензент:

д.ф.-м.н. доцент

Бодунов Николай Александрович

Санкт-Петербург

2022 год

St Petersburg State University

TARASOVA Mariia

Graduation work

A sociological model with nonstandard norming

Level of education: Master

Direction 01.04.01 “Mathematics”

The main educational program BM.5832.2020 “Advanced Mathematics”

Supervisor:

Professor, St Petersburg State University,
Doctor of mathematics, professor
Pilyugin Sergei Yurievich

Reviewer:

Chair of the Department of Mathematics,
St Petersburg State Electrotechnical
University (LETI),
Doctor of mathematics, docent
Bodunov Nikolai Alexandrovich

St Petersburg
2022

1 Introduction

Whenever there is a group of people sharing different opinions and interacting with each other, there is a question about the dynamics of their opinions. Will they reach the consensus? Or will their opinions polarize?

Various models of opinion dynamics have been studied since 1950s ([7, 8]). At present, opinion studies are a well-developed field of research (see, for example, the monographs [15, 20] and the recent survey [18]). The main goal of opinion dynamics is to describe and analyze evolution of public opinion in social systems.

When trying to describe the opinion dynamics problem mathematically, we should set what is the opinion and how the interaction between the people looks like. The opinion spaces can vary a lot. They can be discrete or continuous, one-dimensional or multidimensional. However, mostly (and in the current text), the models of opinion spaces as intervals in \mathbb{R} . And the ways of interaction can be very diverse and complicated.

Mostly, models studied in opinion dynamics are linear. DeGroot [5] model is a simple example of such a model. In this system the vector of opinions an time n is obtained from the previous one by multiplying by a row stochastic matrix. In this case, the entries of the matrix determine how much an agent influences the opinion of another one. This interaction model does not depend on time, so the result opinion of an agent is affected only by the degree of impact of his/her friend on him/her and the degree of the agent's own self-confidence. The simplicity of this model allows one to apply more or less standard methods of linear dynamical systems and describe the dynamics precisely.

One of the first nonlinear models was suggested in [12, 13], where the notion of “bounded confidence” had been introduced. This notion formalizes the fact that, in the course of formation of public opinion, a member of the society is mostly influenced by individuals sharing a similar opinion. Apart from DeGroot model, in this case the group of people who influence the opinion of the given agent can change over time. This can (and does) lead to the significant changes in the dynamics.

The first opinion model based on the notion of bounded confidence, introduced by Hegselmann and Krause, was later called the Hegselmann – Krause (HK) model; this model and its generalizations have been intensively studied by various authors, see, for example, [6, 17, 14, 1, 16, 4, 3, 11, 21, 22, 9]. Mostly, the results were based on computer simulations, and it was noted that “rigorous analytical results are difficult to obtain” [10].

In the paper [18], a modification of the HK model suggested by Campi was studied. Let us consider the dynamics of opinions in a society of voters who have to choose between two options, -1 and 1. Assume that the society is formed by N individuals, and let $v_k^n \in [-1, 1]$ be the opinion of individual k at time moment n .

Fix a positive $\varepsilon < 1$ (the level of bounded confidence in the society) and consider for $k \in \{1, \dots, N\}$ the set of indices

$$J(v_k^n) = \{l \in \{1, \dots, N\} : |v_l^n - v_k^n| \leq \varepsilon\}.$$

This is the set of indices of agents whose opinions influence agent k at time moment n .

In the classical HK model, the dynamics of voters is based on the following procedure: At the step of the process of opinion formation at time n , the new opinion of agent k is obtained by adding to v_k^n a value proportional to the average of values $v_l^n - v_k^n$ over indices l belonging to the set $J(v_k^n)$.

In the model studied in the paper [18], the average of values $v_l^n - v_k^n$ over the set $J(v_k^n)$ is replaced by the average of values v_l^n over the set $J(v_k^n)$.

Thus, when taking the average, the opinion of agent k is included into consideration. In the case where an agent has no other agents with ε -close opinions, this means that the agent enforces her/his belief: In absence of counter-arguments, one tends to strengthen her/his own opinion.

Mathematically, this modification of the process may lead to the following consequence: some of the new values may be outside the interval $[-1, 1]$. In [18], a “cutting” procedure was suggested; the new opinion value is obtained by replacing the values less than -1 by -1, and the values more than 1 are replaced by 1.

The dynamics of the appearing dynamical system had been completely described in [18]. It was shown that if $\varepsilon \leq 1/2$, then any trajectory tends to a fixed point as time goes to infinity. All possible fixed points had been characterized. It was shown that any fixed point $P = (p_1, \dots, p_N)$ with $|p_k| = 1$, $k \in \{1, \dots, N\}$, is attracting, while all the remaining fixed points are Lyapunov unstable. Modifications of the model studied in [18] were considered in the recent papers [2] (where the average of values v_l^n has been replaced by values $i(v_l^n)$ for a wide class of influence functions i) and [19] (where the finite set of agents has been replaced by the continuum $[0, 1]$).

In this thesis, we study a model similar to that considered in [18] but with a different norming. Instead of “cutting” the values obtained at the first step (when we add to v_k^n a value proportional to the average of values v_l^n over the set $J(v_k^n)$ and obtain values w_k^n), now we divide the values w_k^n by the maximal absolute value of w_k^n (see a detailed description of the appearing dynamical system in the next section).

Our main results are as follows:

- we find a condition (see inequality (6)) under which any trajectory tends to a fixed point as time goes to infinity;
- we describe all fixed points in this case;
- we show that both fixed points $P_- = (-1, \dots, -1)$ and $P_+ = (1, \dots, 1)$ are attracting;
- we prove that all the remaining fixed points are Lyapunov stable (thus, the dynamics of our system is completely different from that of the system studied in [18]) but not attracting;
- we give an example of the system for which condition (6) is not satisfied and that has an unstable fixed point.

Of course, our reasoning in this text essentially differs from that in [18].

The structure of the thesis is as follows. Section 2 is devoted to the statement of the problem. In Section 3, basic properties of the system are described. In Section 4, we prove the convergence of trajectories to fixed points. In Section 5, stability of fixed points is analyzed. In Section 6, we give an example of a system with an unstable fixed point. Section 7 contains several examples of computer modeling.

2 Statement of the problem

We study a dynamical system modeling the following problem of opinion dynamics. A society consisting of N agents has to choose between two options, 1 and -1. Let $v_k^n \in [-1, 1]$ be the opinion of agent with index $k \in \{1, \dots, N\}$ at time moment $n = 0, 1, \dots$ and let

$$V^n = (v_1^n, \dots, v_N^n)$$

be the array of opinions at time moment n .

Let us define the operator Φ determining the iterative process which models the opinion dynamics. Fix two numbers $h, \varepsilon \in (0, 1)$ and an array

$$V = (v_k \in [-1, 1] : k \in \{1, \dots, N\}).$$

Introduce the sets

$$J(v_k) = \{l \in \{1, \dots, N\} : |v_l - v_k| \leq \varepsilon\}, \quad k \in \{1, \dots, N\}.$$

Denote by $I(v_k)$ the cardinality of the (nonempty) set $J(v_k)$.

Define an auxiliary array

$$W(V) = (w_1(V), \dots, w_N(V)),$$

where

$$w_k(V) = v_k + \frac{h}{I(v_k)} \sum_{l \in J(v_k)} v_l, \quad k = 1, \dots, N. \quad (1)$$

Now, assuming that $W(V)$ is a nonzero array, we set

$$m(W(V)) = \max_{l \in \{1, \dots, N\}} |w_l(V)|$$

and

$$\Phi(V) = (v'_1, \dots, v'_N),$$

where

$$v'_k = \frac{w_k(V)}{m(W(V))}.$$

Clearly,

$$v'_k \in [-1, 1]. \quad (2)$$

We can represent $W(V)$ in the form

$$W(V) = (E + hA)V,$$

where E is the identity matrix and the matrix A is row-stochastic; it easily follows from the inclusion $h \in (0, 1)$ that if $V \neq 0$, then $W(V) \neq 0$ as well.

Consider an initial array of opinions $V^0 = (v_1^0, \dots, v_N^0)$. If $V^0 = 0$, then we set $V^n = 0$ for $n \geq 0$ and exclude this trivial case from the further consideration.

It follows from the above reasoning that if $V^0 \neq 0$, then $V^n = \Phi^n(V^0)$ are defined for $n > 0$. Our main goal is to study the behavior of positive trajectories of the appearing dynamical system.

3 Basic properties of the system

For simplicity, we denote $w_k^n = w_k(V^n)$.

Denote by \mathcal{V} the set of arrays V such that

$$v_1 \leq v_2 \leq \dots \leq v_N. \quad (3)$$

Lemma 3.1. *If $v \in \mathcal{V}$, then $\Phi(V) \in \mathcal{V}$.*

Proof. It follows from [18, Corollary 1] that inequality (3) implies the inequalities $w_k(V) \leq w_{k+1}(V)$; division by $m(W(V))$ preserves the required inequalities. \square

It is easily seen that the value $w_k(V)$ in (1) does not depend on the indexing of components of V . Hence, in what follows, we may (and will) consider trajectories belonging to \mathcal{V} . \square

Let us note one important inequality. Without loss of generality, we may assume that, for given V , $m(W(V)) = |w_N(V)|$. Then

$$m(W(V)) = \left| v_N + \frac{h}{I(v_N)} \sum_{l \in J(v_N)} v_l \right| \leq 1 + h. \quad (4)$$

Lemma 3.2. *If $|v_k^n - v_{k+1}^n| > \varepsilon$, then $|v_k^\nu - v_{k+1}^\nu| > \varepsilon$ for all $\nu > n$.*

Proof. It is enough to prove the statement for $\nu = n + 1$.

The inequality $|v_k^n - v_{k+1}^n| > \varepsilon$ implies that $J(v_k^n) \subset \{1, \dots, k\}$ and $J(v_{k+1}^n) \subset \{k + 1, \dots, N\}$. Hence,

$$v_k^n + \frac{h}{I(v_k^n)} \sum_{l \in J(v_k^n)} v_l^n \leq v_k^n(1 + h)$$

and

$$v_{k+1}^n + \frac{h}{I(v_{k+1}^n)} \sum_{l \in J(v_{k+1}^n)} v_l^n \geq v_{k+1}^n (1+h).$$

Thus,

$$v_{k+1}^{n+1} - v_k^{n+1} \geq \frac{(v_{k+1}^n - v_k^n) \cdot (1+h)}{m(W(V^n))} > \frac{\varepsilon(1+h)}{m(W(V^n))}. \quad (5)$$

Now inequality (4) implies the required inequality

$$v_{k+1}^{n+1} - v_k^{n+1} > \varepsilon.$$

□

Remark 1. An analog of Lemma 3.2 does not necessarily hold for v_k^n and v_m^n with $|k-m| \neq 1$. Let us consider the following example.

Let $N = 8$, $\varepsilon = 1/2$, and $h = 1/3$. If

$$V^0 = (-1, -5/16, 0, 0, 0, 0, 5/16, 1),$$

then $m(W(V^0)) = 1+h$ and

$$V^1 = (-1, -1/4, 0, 0, 0, 0, 1/4, 1).$$

Hence, $v_7^0 - v_2^0 = 10/16 > \varepsilon$, while $v_7^1 - v_2^1 = 1/2 = \varepsilon$.

4 Convergence to fixed points

In this section, we show that if

$$\varepsilon(N-1) < 1, \quad (6)$$

then any trajectory $\Phi^n(V)$ converges to a fixed point of Φ as $n \rightarrow \infty$.

Now we introduce the object which is the main tool in the following proofs.

Definition 4.1. For an array $V^n = \Phi^n(V^0)$, a set $\{k, k+1, \dots, m\} \subset \{1, \dots, N\}$ is called a *band at time n* if the following properties are satisfied:

- (1) if $k > 1$, then $|v_{k-1}^n - v_k^n| > \varepsilon$, and if $m < N$, then $|v_m^n - v_{m+1}^n| > \varepsilon$;
- (2) $|v_l^n - v_{l+1}^n| \leq \varepsilon$ for all $l \in \{k, \dots, m-1\}$.

The value $|v_k^n - v_m^n|$ is called the *diameter* of the band $\{k, k+1, \dots, m\}$.

In what follows, we often use the term *band* instead of *band at time n*.

Remark 2. Since we work with trajectories in \mathcal{V} ,

$$v_k^n \leq \dots \leq v_m^n$$

for any band $\{k, k+1, \dots, m\}$.

It follows from Lemma 3.2 that if $\{k, k + 1, \dots, m\}$ is a band at time n for some $V^n = \Phi^n(V^0)$, then no subset of $\{k, k + 1, \dots, m\}$ can be a subset of a band at time $\nu > n$ for V^ν containing either $k - 1$ or $m + 1$. Hence, either a band $\{k, k + 1, \dots, m\}$ at time n for V^n is a band at time $n + 1$ for V^{n+1} as well or it splits into a union of several bands of smaller lengths.

Thus, for any band $\{k, k + 1, \dots, m\}$ of any initial array V^0 there exists a unique decomposition

$$\{k, k + 1, \dots, m\} = \bigcup_{j=1}^r \{k_j, \dots, m_j\} \quad (7)$$

with $k_1 = k$, $m_r = m$, and $k_{j+1} = m_j + 1$ and a time ν such that any $\{k_j, \dots, m_j\}$ is a band for any $V^n = \Phi^n(V^0)$ for any $n \geq \nu$ (i.e., it does not split into bands of smaller lengths as time grows).

Clearly, if V^0 is a nonzero array, then either $v_1^n = -1$ or $v_N^n = 1$ for any $n > 0$. We assume that the same holds for $n = 0$.

We introduce the following condition on the initial array V^0 .

Condition A. *The array V^0 has a band $\{k, \dots, m\}$ at time 0 such that v_k^0, \dots, v_m^0 are nonzero and have the same sign.*

If V^0 contains a single band, then this band is $\{1, \dots, N\}$, and, by our assumption, either $v_1^0 = -1$ either $v_N^0 = 1$. Then inequality (6) implies that V^0 satisfies Condition A.

If V^0 contains at least two bands, then it obviously satisfies Condition A.

Thus, inequality (6) implies Condition A for any array V^0 .

In the remaining part of this section, we assume that any initial array V^0 satisfies Condition A.

Lemma 4.2. *For any V^0 , the following relation holds:*

$$\lim_{n \rightarrow \infty} m(W(V^n)) = 1 + h. \quad (8)$$

Proof. Fix a band $\{k, \dots, m\}$ for V^0 at time 0 such that v_k^0, \dots, v_m^0 are nonzero and have the same sign. Without loss of generality, we may assume that these values are positive.

Let us consider the behavior of v_k^n as n grows:

$$v_k^{n+1} = \frac{1}{m(W(V^n))} \left(v_k^n + h \frac{1}{I(v_k^n)} \sum_{l \in J(v_k^n)} v_l^n \right) \geq \frac{(1+h)}{m(W(V^n))} v_k^n.$$

To get a contradiction, assume that relation (8) does not hold. Then there exists a subsequence n_k tending to infinity such that

$$m(W(V^{n_k})) \leq 1 + h - \alpha$$

for some $\alpha \in (0, h)$.

Without loss of generality, we may assume that the above inequalities hold for all n . Then

$$v_k^{n+1} \geq \frac{1+h}{1+h-\alpha} v_k^n = \beta v_k^n,$$

where

$$\beta = \frac{1+h}{1+h-\alpha} > 1.$$

Thus,

$$v_k^{n+1} \geq \beta^n v_k^0 \rightarrow \infty, \quad n \rightarrow \infty,$$

which contradicts the inequalities $v_k^n \leq 1$. \square

Let us describe the behavior of a band of diameter not more than ε .

Lemma 4.3. *Assume that a band $\{k, \dots, m\}$ for an array V^ν at time ν has diameter not more than ε and does not split as time grows. Then the diameters of this band for the arrays V^n at all times $n > \nu$ are not more than ε as well and*

$$\lim_{n \rightarrow \infty} (v_m^n - v_k^n) = 0. \quad (9)$$

Proof. Without loss of generality, we assume that $\nu = 0$. Applying Lemma 4.2, we may also assume that

$$m(W(V^n)) \geq \gamma > 1, \quad n \geq 0.$$

It follows from our assumption (the band does not split) that $J(v_k^n) = J$ and $I(v_k^n) = I$ for $n \geq 0$. Hence, if $n \geq 0$, then

$$\begin{aligned} v_m^{n+1} - v_k^{n+1} &= \frac{1}{m(W(V^{n+1}))} \left(v_m^n + \frac{h}{I} \sum_{l \in J} v_l^n - v_k^n - \frac{h}{I} \sum_{l \in J} v_l^n \right) = \\ &= \frac{1}{m(W(V^{n+1}))} (v_m^n - v_k^n) \leq \frac{1}{\gamma} (v_m^n - v_k^n). \end{aligned}$$

This obviously implies the statement of our lemma. \square

Now we prove that the diameter of every band in decomposition (7) does not exceed ε for times $n \geq \nu$ if ν is large enough.

Lemma 4.4. *For any initial array V^0 and any its band $\{k, \dots, m\}$ at time 0 there exists a time ν such that the diameter of any band $\{k_j, \dots, m_j\}$ in decomposition (7) for V^n with $n \geq \nu$ does not exceed ε .*

First we fix some constants.

Fix a positive δ such that

$$\delta \leq \frac{\varepsilon h}{3N(1+h)}, \quad (10)$$

a positive β such that

$$\beta \leq \frac{\delta h}{3N(1+h)}, \quad (11)$$

and a positive α such that

$$\frac{\alpha}{1+h-\alpha} \leq \beta. \quad (12)$$

We get Lemma 4.4 as a corollary of the following two lemmas.

Lemma 4.5. *Assume that the diameter of a band $\{k, \dots, m\}$ is larger than ε for all $n \geq 0$. There exists an $n_0 \geq 0$ such that the following implication holds. If $v_l^n \in [v_k^n, v_k^n + \delta)$ for all $l \in J(v_k^n)$ and for some $n \geq n_0$, then there exists an index s such that $v_s^{n+1} \in [v_k^{n+1} + \delta, v_k^{n+1} + \varepsilon]$.*

Proof. Fix a time n and let $s = \max J(v_k^n)$. We will show that the assumption of the lemma implies the inclusion $v_s^{n+1} \in [v_k^{n+1} + \delta, v_k^{n+1} + \varepsilon]$.

By assumption, the value $v_m^n - v_k^n$ (the diameter of the band $\{k, \dots, m\}$) is larger than ε for all n , and it follows from the inequality

$$v_s^n - v_k^n \leq \varepsilon$$

that $s < m$. Hence, $s, s+1 \in \{k, \dots, m\}$. Since $\{k, \dots, m\}$ is a band, $s+1 \in J(v_s^n)$.

Then we have the following estimates for $n \geq 0$:

$$\begin{aligned} v_s^{n+1} - v_k^{n+1} &= \frac{1}{m(W(V^n))} \left(v_s^n + \frac{h}{I(v_s^n)} \sum_{l \in J(v_s^n)} v_l^n - v_k^n - \frac{h}{I(v_k^n)} \sum_{l \in J(v_k^n)} v_l^n \right) \geq \\ &\geq \frac{1}{m(W(V^n))} \left(v_s^n + h \left(\frac{I(v_s^n) - 1}{I(v_s^n)} v_k^n + \frac{1}{I(v_s^n)} v_{s+1}^n \right) - v_k^n - h v_s^n \right) \geq \\ &\geq \frac{1}{m(W(V^n))} \left(v_s^n + h \left(\frac{I(v_s^n) - 1}{I(v_s^n)} (v_s^n - \delta) + \right. \right. \\ &\quad \left. \left. + \frac{1}{I(v_s^n)} (v_s^n + \varepsilon - \delta) \right) - v_k^n - h(v_k^n + \delta) \right) \geq \\ &\geq \frac{1+h}{m(W(V^n))} (v_s^n - v_k^n) + \frac{h}{m(W(V^n))} \cdot \left(\frac{\varepsilon}{N} - 2\delta \right) \geq \\ &\geq \frac{h}{1+h} \left(\frac{\varepsilon}{N} - 2\delta \right) \geq \delta. \end{aligned}$$

In the last line, we apply inequality (10).

To get the upper bound, take a number $n_0 \geq 0$ such that

$$m(W(V^n)) \geq \frac{2}{3} + h, \quad n \geq n_0.$$

If $n \geq n_0$, then

$$\begin{aligned}
v_s^{n+1} - v_k^{n+1} &= \frac{1}{m(W(V^n))} \left(v_s^n + \frac{h}{I(v_s^n)} \sum_{l \in J(v_s^n)} v_l^n - v_k^n - \frac{h}{I(v_k^n)} \sum_{l \in J(v_k^n)} v_l^n \right) \leq \\
&\leq \frac{1}{m(W(V^n))} \left(v_s^n + h \left(\frac{1}{I(v_s^n)} v_k^n + \frac{I(v_s^n) - 1}{I(v_s^n)} v_{s+1}^n \right) - v_k^n - h v_k^n \right) \leq \\
&\leq \frac{1}{m(W(V^n))} \left(v_k^n + \delta + h \left(\frac{1}{I(v_s^n)} v_k^n + \frac{I(v_s^n) - 1}{I(v_s^n)} (v_k^n + \varepsilon + \delta) \right) - v_k^n - h v_k^n \right) = \\
&= \frac{1}{m(W(V^n))} \left(\delta + h \left(\frac{I(v_s^n) - 1}{I(v_s^n)} (\varepsilon + \delta) \right) \right) \leq \\
&\leq \frac{1}{\frac{2}{3} + h} ((1 + h)\delta + h\varepsilon) \leq \frac{1}{\frac{2}{3} + h} (2\delta + h\varepsilon) \leq \varepsilon,
\end{aligned}$$

where in the last line we take into account that $\delta \leq \frac{\varepsilon}{3}$ due to (10). \square

Lemma 4.6. *There exists an $n_0 \geq 0$ such that for any $n \geq n_0$, the following inequality holds:*

$$v_k^{n+2} \geq v_k^n + \beta.$$

Proof. We claim that there exists an n_0 such that

$$v_k^{n+1} \geq v_k^n - \beta, \quad n \geq n_0, \quad (13)$$

and if there exists an index s such that $v_s^n \in [v_k^n + \delta, v_k^n + \varepsilon]$, then

$$v_k^{n+1} \geq v_k^n + 2\beta, \quad n \geq n_0. \quad (14)$$

Then the statement of our lemma follows from Lemma 4.5. To establish estimate (13), let us fix a positive α such that inequality (12) holds.

Consider an $n_0 \geq 0$ such that

$$m(W(V^n)) \geq 1 + h - \alpha, \quad n \geq n_0.$$

If $n \geq n_0$, then

$$\begin{aligned}
v_k^{n+1} &= \frac{1}{m(W(V^n))} \left(v_k^n + \frac{h}{I(v_k^n)} \sum_{l \in J(v_k^n)} v_l^n \right) \geq \frac{1 + h}{m(W(V^n))} v_k^n = \\
&= v_k^n + \left(\frac{1 + h}{m(W(V^n))} - 1 \right) v_k^n \geq v_k^n - \left(\frac{1 + h}{m(W(V^n))} - 1 \right) \geq \\
&\geq v_k^n - \frac{\alpha}{1 + h - \alpha} \geq v_k^n - \beta.
\end{aligned}$$

Next, we assume that $v_s^n \in [v_k^n + \delta, v_k^n + \varepsilon]$. Let us estimate

$$\begin{aligned}
v_k^{n+1} &= \frac{1}{m(W(V^n))} \left(v_k^n + \frac{h}{I(v_k^n)} \sum_{l \in J(v_k^n)} v_l^n \right) \geq \\
&\geq \frac{1}{m(W(V^n))} \left(v_k^n + h \left(\frac{I(v_k^n) - 1}{I(v_k^n)} v_k^n + \frac{1}{I(v_k^n)} v_s^n \right) \right) \geq \\
&\geq \frac{1}{m(W(V^n))} \left(v_k^n + h \left(\frac{I(v_k^n) - 1}{I(v_k^n)} v_k^n + \frac{1}{I(v_k^n)} (v_k^n + \delta) \right) \right) = \\
&= \frac{1+h}{m(W(V^n))} v_k^n + \frac{\delta h}{I(v_k^n) \cdot m(W(V^n))} \geq \\
&\geq v_k^n - \frac{\alpha}{1+h-\alpha} + \frac{\delta h}{N(1+h)} \geq v_k^n + 2\beta.
\end{aligned}$$

□

To prove Lemma 4.4, we assume that the diameter of the band $\{k, \dots, m\}$ is more than ε for arbitrarily large n_0 . Then Lemmas 4.5 and 4.6 lead to a contradiction since the sequence (v_k^n) is bounded.

Theorem 4.7. *If condition (6) is satisfied, then any trajectory $\Phi^n(V^0)$ tends to a fixed point of Φ as $n \rightarrow \infty$.*

Proof. It follows from Lemmas 4.3 and 4.4 that for any initial nonzero array V^0 the following holds: if n is large enough, then the set $\{1, \dots, N\}$ is the union of disjoint subsets,

$$\{1, \dots, N\} = \bigcup_{i=1}^r \{k : b_i \leq k \leq c_i\},$$

where $b_1 = 1$, $c_r = N$, $c_{i+1} = b_i + 1$, and any set $\{b_i, \dots, c_i\}$ is a band for V^n such that

$$0 \leq v_{c_i}^n - v_{b_i}^n \rightarrow 0, \quad n \rightarrow \infty.$$

There are the following possible cases:

- $v_{b_i}^n \geq 0$ for some n ; then $v_k^m \geq 0$ for all $b_i \leq k \leq c_i$ and $m \geq n$;
- $v_{c_i}^n \leq 0$ for some n ; then $v_k^m \leq 0$ for all $b_i \leq k \leq c_i$ and $m \geq n$;
- $v_{b_i}^n v_{c_i}^n < 0$ for some n ; then $v_k^m v_k^m < 0$ for all $b_i \leq k \leq c_i$ and $m \geq n$.

In any of these cases, there exist numbers $a_i \in [0, 1]$ such that

$$v_k^n \rightarrow a_i, \quad b_i \leq k \leq c_i, \quad n \rightarrow \infty,$$

(and $a_i = 0$ in the third case).

It follows from the left-hand side of inequality (5) that

$$v_{c_{i+1}}^{n+1} - v_{b_i}^{n+1} \geq v_{c_{i+1}}^n - v_{b_i}^n;$$

hence,

$$a_{i+1} - a_i > \varepsilon, \quad i = 1, \dots, r-1.$$

In addition, either $a_1 = -1$ or $a_r = 1$ (or both possibilities are realized). Clearly, the corresponding array

$$A = (a_1, \dots, a_1, a_2, \dots, a_2, \dots, a_r, \dots, a_r)$$

is a fixed point of Φ such that

$$V^n \rightarrow A, \quad n \rightarrow \infty. \quad \square$$

Remark 3. In fact, the proofs of Lemmas 4.3 and 4.4 (and hence, of Theorem 4.7) are based not on condition (6) but on the assumption that for any V^0 , relation (8) holds (which we deduce from Condition A).

We refer to condition (6) in Theorem 4.7 since the above-formulated two assumptions are of “inner” character while condition (6) relates values from the statement of the problem.

5 Stability of fixed points

In this section, we assume that relation (8) holds. As was noted, this assumption implies the conclusion of Theorem 4.7. Let us study the stability properties of the appearing fixed points of Φ .

First we introduce the following notation. Let $P = (p_1, \dots, p_N)$ be a fixed point of Φ . An array

$$(B_1(a_1), \dots, B_r(a_r))$$

is called the *scheme* of the fixed point P if any

$$B_j(a_j) = \{b_j, \dots, c_j\}, \quad j = 1, \dots, r,$$

where $b_1 = 1$, $c_r = N$, and $c_{i+1} = b_i + 1$, is a band for $P = \Phi^n(P)$ at any time n and

$$p_k = a_j, \quad k \in B_j(a_j).$$

Let us select two fixed points, P_- and P_+ , having schemes $(B_1(-1))$ and $(B_1(1))$, respectively, where $B_1(-1) = B_1(1) = \{1, \dots, N\}$.

Theorem 5.1. *If relation (8) holds, then*

- (1) *both fixed points P_- and P_+ are asymptotically stable for Φ ;*
- (2) *any fixed point P different from P_- and P_+ is Lyapunov stable but not asymptotically stable.*

Proof. Let us first prove item (1). We consider the case of the fixed point P_+ , for P_- the proof is similar.

First we prove that P_+ is Lyapunov stable. Fix a $\Delta > 0$; without loss of generality, we assume that $\Delta \leq \varepsilon$.

Let $\delta \leq \Delta$ and consider a $V = (v_1, \dots, v_N) \in \mathcal{V}$ such that

$$v_k \in [1 - \delta, 1], \quad k \in \{1, \dots, N\}. \quad (15)$$

Then $J(v_k) = \{1, \dots, N\}$ for $k \in \{1, \dots, N\}$; hence,

$$w_k(V) \geq (1 + h)(1 - \delta), \quad k \in \{1, \dots, N\},$$

and

$$(\Phi(V))_k \in [1 - \delta, 1], \quad k \in \{1, \dots, N\}.$$

This implies that

$$(\Phi^n(V))_k \in [1 - \Delta, 1], \quad k \in \{1, \dots, N\}, \quad n \geq 0.$$

Thus, P_+ is Lyapunov stable.

As was said before introducing Condition A, we may assume that, for any V , either $(\Phi^n(V))_1 = -1$ or $(\Phi^n(V))_N = 1$ for $n \geq 0$. In our case, inequality (15) implies that $v_N^n = (\Phi^n(V))_N = 1$ for all n , and it follows from Lemma 4.3 that

$$|v_k^n - 1| \rightarrow 0, \quad n \rightarrow \infty, \quad k \in \{1, \dots, N\}.$$

Thus, P_+ is asymptotically stable.

Now we prove item (2). Consider two possible cases.

Case 1. The fixed point P has scheme $(B_1(-1), B_2(1))$ with nonempty $B_1(-1), B_2(1)$. In this case, the Lyapunov stability is proved by the same reasoning as above. To prove that P is not asymptotically stable, note that any point with scheme $(B_1(-1 + \delta), B_2(1))$, where $\delta \in (0, \varepsilon)$, is a fixed point of Φ .

Case 2. The fixed point $P = (p_1, \dots, p_n)$ has scheme

$$(B_1(-1), \dots, B_l(a_l), \dots, B_r(1)),$$

where $B_l(a_l)$ is nonempty and $|a_l| \neq 1$. In this case, $|a_l| \in (-1 + \varepsilon, 1 - \varepsilon)$ and one of the sets $B_1(-1), B_r(1)$ is nonempty. To simplify consideration, assume that $B_1(-1)$ is empty (the remaining cases are treated similarly).

Fix a $\Delta > 0$ such that

$$a_{j+1} - a_j > \varepsilon + 2\Delta, \quad j = 1, \dots, r - 1. \quad (16)$$

Without loss of generality, we assume that $\Delta \leq \varepsilon/2$. Clearly, if $V = (v_1, \dots, v_N)$ and

$$|v_k - p_k| \leq \Delta, \quad k \in \{1, \dots, N\},$$

then $J(v_k) = J(p_k)$ for $k \in \{1, \dots, N\}$.

Take a positive δ such that

$$\frac{2\delta}{1-\delta} < \Delta. \quad (17)$$

Clearly, in this case $\delta < \Delta$.

Introduce the following condition on the trajectory of an initial point V^0 .

Condition $C(\nu)$:

$$|v_k^n - p_k| \leq \Delta, \quad k \in \{1, \dots, N\}, \quad 0 \leq n \leq \nu.$$

We show that if

$$|v_k^0 - p_k| \leq \delta, \quad k \in \{1, \dots, N\}, \quad (18)$$

for an initial point V^0 , then Condition $C(\nu)$ is satisfied for all $\nu \geq 0$, which, of course, means that P is Lyapunov stable.

Thus, below we assume that inequalities (18) are satisfied.

Since $\delta < \Delta$, Condition $C(0)$ is satisfied. Now we show that Condition $C(\nu)$ implies Condition $C(\nu + 1)$.

We start with $k \in B_r(1)$. Due to Condition $C(\nu)$, $J(v_k^n) = B_r(1)$ for $0 \leq n \leq \nu$. The same reasoning as in the proof of item (1) shows that

$$|v_k^n - 1| < \delta, \quad k \in B_r(1), \quad n \leq \nu + 1.$$

Denote

$$\mu_n = m(W(V^{n-1})) \times \dots \times m(W(V^1)) \times m(W(V^0)).$$

Since $v_k^0 \geq 1 - \delta$ for $k \in B_r(1)$,

$$\frac{(1+h)^n}{\mu_n}(1-\delta) \leq v_k^n \leq 1, \quad 0 \leq n \leq \nu + 1, \quad k \in B_r(1),$$

and the inequalities

$$\frac{(1+h)^n}{\mu_n} - 1 \leq \frac{\delta}{1-\delta}, \quad 0 \leq n \leq \nu + 1, \quad (19)$$

hold.

Now we consider indices $k \in B_l(a_l)$ with $l < r$. Condition $C(\nu)$ and inequalities (18) imply that

$$v_k^0 \leq a_l + \delta, \quad v_k^1 \leq \frac{1+h}{\mu_1}(a_l + \delta), \quad \dots, \quad v_k^n \leq \frac{(1+h)^n}{\mu_n}(a_l + \delta)$$

for $0 \leq n \leq \nu + 1$.

Hence, if $k \in B_l(a_l)$, then it follows from inequality (19) with $n = \nu + 1$ that

$$\begin{aligned} v_k^{\nu+1} - a_l &\leq \frac{(1+h)^{\nu+1}}{\mu_{\nu+1}}(a_l + \delta) - a_l = \\ &= \left(\frac{(1+h)^{\nu+1}}{\mu_{\nu+1}} - 1 \right) a_l + \frac{(1+h)^{\nu+1}}{\mu_{\nu+1}} \delta \leq \frac{2\delta}{1-\delta} < \Delta. \end{aligned}$$

Similarly one shows that

$$v_k^{\nu+1} - a_l > -\Delta,$$

which proves that Condition C($\nu + 1$) is satisfied.

This completes the proof of Lyapunov stability of the fixed point P .

To prove that P is not asymptotically stable, note that any point with scheme

$$(B_1(-1), \dots, B_l(a_l + \delta), \dots, B_r(1)),$$

where δ is small enough, is a fixed point of Φ . □

6 Example with a single band

The following example shows that if ε is not small, then the dynamics of the system can be essentially different from that described above.

Let $N = 6$ and $\varepsilon = 1/2$. Then Φ has a fixed point

$$P = (-1, -1/2, 0, 0, 1/2, 1).$$

Clearly,

$$\begin{aligned} W(P) &= \left(-1 - \frac{3h}{4}, -\frac{1}{2} - \frac{3h}{8}, 0, 0, \frac{1}{2} + \frac{3h}{8}, 1 + \frac{3h}{4} \right), \\ m(W(P)) &= 1 + \frac{3h}{4}, \end{aligned} \tag{20}$$

and $\Phi(P) = P$.

This fixed point is unstable; for any small $\delta > 0$, the point

$$V^0 = (-1, -1/2, \delta, \delta, 1/2, 1)$$

has a band $\{\delta, \delta, 1/2, 1\}$ at time 0 with nonzero elements of the same sign; the reasoning applied in the proof of Lemma 4.2 shows that

$$m(W(V^n)) \rightarrow 1 + h, \quad n \rightarrow \infty,$$

which, compared with relation (20) indicates that the fixed point P is unstable.

In Fig. 3 of the next section, the dynamics with $\delta = 0.01$ is shown.

7 Numerical experiments

The first figure illustrates the dynamics of the system for which condition (6) holds. Figure 1 shows the initial distribution of opinions and the evolution of the system at times 10, 30, and 70. One can see that at time 70, the equilibrium is almost reached. The fixed points of this system are 4 groups of equal numbers.




Figure 1: Initial distribution and opinions' evolution of system with (6) at steps 10, 30 and 70; $\varepsilon = 0.1$, $h = 0.1$.




Figure 2: Initial distribution and opinions' evolution of system with Condition A at steps 20, 40 and 90; $\varepsilon = 0.4$, $h = 0.1$.




Figure 3: Initial distribution and opinions' evolution for third example at steps 10, 30 and 70, when the equilibrium is reached; $\varepsilon = 0.5$, $h = 0.1$.

The second example illustrates the evolution of the system with a larger number of agents. Here, condition (6) is not met, but Condition A holds. Figure 2 shows the initial distribution of such a system with $\varepsilon = 0.5$, $h = 0.1$ and its evolution at times 20, 40, and 90.

Figure 3 shows the dynamics of the band

$$V^0 = (-1, -1/2, \delta, \delta, 1/2, 1)$$

which was mentioned in Section 6. The illustration shows the initial distribution of this system with $\delta = 0.01$ and its evolution at times 10, 30, and 70. In this case, the band of positive values collapses into a band with the same values.

References

- [1] M. Bertotti and M. Delitala, Cluster formation in opinion dynamics: A qualitative analysis, *Z. Angew. Math. Phys.*, **61** (2010), 583–602.
- [2] N. A. Bodunov and S. Y. Pilyugin, Convergence to Fixed Points in One Model of Opinion Dynamics, *J. Dyn. Control. Syst.*, **27** (2021), 617–623.
- [3] D. Borra and T. Lorenzi, Asymptotic analysis of continuous opinion dynamics models under bounded confidence *Commun. Pure Appl. Anal.*, **12** (2013), 1487–1499.
- [4] F. Ceragioli and P. Frasca, Continuous and discontinuous opinion dynamics with bounded confidence, *Nonlinear Anal. Real World Appl.*, **13** (2012), 1239–1251.
- [5] M. DeGroot, Reaching a consensus, *Journal of the American Statistical Association*, **69** (1974), 118–121.
- [6] S. Fortunato, The Krause – Hegselmann consensus model with discrete opinions, *Internat. J. Modern Phys. C*, **15** (2004), 1021–1029.
- [7] J. French, A formal theory of social power, *Psychological Review*, **63** (1956), 181–194.
- [8] F. Harary, A criterion for unanimity in French’s theory of social power, in *Studies in Social Power* (ed. D. Cartwright), Institute for Social Research, Ann Arbor, 1959.
- [9] P. Hegarty and E. Wedin, The Hegselmann – Krause dynamics for equally spaced agents, *J. Difference Equ. Appl.*, **22** (2016), 1621–1645.
- [10] R. Hegselmann and U. Krause, Opinion dynamics and bounded confidence: Models, analysis and simulation, *Journal of Artificial Societies and Social Simulation*, **5** (2002), 1–33.
- [11] P. Jabin and S. Motsch, Clustering and asymptotic behavior in opinion formation, *J. Differential Equations*, **257** (2014), 4165–4187.
- [12] U. Krause, A discrete nonlinear and non - autonomous model of consensus formation, in *Communications in Difference Equations* (ed. S. E. et al.), Gordon and Breach Publ., Amsterdam, 1997.
- [13] U. Krause, Soziale dynamiken mit vielen interakteuren. eine problemskizze, in *Modellierung und Simulation von Dynamiken mit vielen interagierenden Akteuren* (eds. U. Krause and M. Stockler), Universitat Bremen, 1997.
- [14] U. Krause, Compromise, consensus, and the iteration of means, *Elem. Math.*, **64** (2009), 1–8.
- [15] U. Krause, *Positive Dynamical Systems in Discrete Time. Theory, Models, and Applications*, De Gruyter, Berlin, 2015.

- [16] S. Kurz and J. Rambau, On the Hegselmann – Krause conjecture in opinion dynamics, *Journal of Difference Equations and Applications*, **17** (2011), 859–876.
- [17] J. Lorenz, Continuous opinion dynamics under bounded confidence: A survey, *International Journal of Modern Physics C*, **18** (2007), 1819–1838.
- [18] S. Y. Pilyugin and M. C. Campi, Opinion formation in voting processes under bounded confidence, *Networks and Heterogeneous Media*, **14** (2019), 617–632.
- [19] S. Y. Pilyugin and D. Z. Sabirova, Dynamics of a Continual Sociological Model, *Vestnik St. Petersburg University: Mathematics*, **54** (2021), 196–205.
- [20] W. Ren and Y. Cao, *Distributed Coordination of Multi-agent Networks. Emergent Problems, Models, and Issues*, Springer, 2011.
- [21] E. Wedin and P. Hegarty, The Hegselmann – Krause dynamics for the continuous-agent model and a regular opinion function do not always lead to consensus, *IEEE Trans. Automat. Control*, **60** (2015), 2416–2421.
- [22] S. Wongkaew, M. Caponigro and A. Borzi, On the control through leadership of the Hegselmann – Krause opinion formation model, *Mathematical Models and Methods in Applied Sciences*, **25** (2015), 565–585.

The main results of this thesis are included in the paper “A model of voting dynamics under bounded confidence with nonstandard norming” that is submitted to the *Networks and Heterogeneous Media*.